COVID 19 (Pfizer-BioNTech) VACCINE QUESTIONNAIRE AND CONSENT 2021-2022 FORMULA

Name	:				D.O	.B. :	Age:	Sex:	
	L	ast Fi	rst						
Schoo	ol		In	nmigrati	on Status _				
Paren	nt Informatio	<u>n</u>							
Resid	ence Address	(include district) _							
Email Address					P	hone			
Natio	nality								
<u>Tick</u>	(√) the app	ropriate answer	to questic	ons belo	<u>w:</u>				
1.	Is your chi	ld suffering from a	fever today	y?		□ Yes	□ No		
2.	Does your child have any of the following medical conditions?								
	□ Impaired Immunity □ Diabetes □ Cancer □ Chronic Lung Disease □ Chronic Liver Disease								
	☐ Chronic Heart Disease ☐ Sickle Cell Disease ☐ Chronic Renal Disease ☐ Morbid Obesity								
	□ Long term Aspirin Therapy □ Other □ None of the above							e above	
3.	Has your child had any serious allergic reaction to a previous vaccine? ☐ Yes ☐ No								
4.	List allergies								
5.	Has your child had COVID 19?					□ Yes		□ No	
	If Yes, W	nen? Date:							
6.	I verify th	at my child has no	ot had any	vaccine v	within the la	st 4 weeks	□ Yes □	□ No	
9.		d the risks and bene					te reverse side of the for my above-name		
	Client /Parent /Guardian Signature:(Circle as appropriate)					Date:			
					ficial use O		[
	anufacturer	Lot # Exp date FN4075	Dose 10mcg/	Route	Site	Date given	Given by (Sign)	Remarks	

IM

0.2 ml

Cayman Islands Public Health Department

03/4/22

Pfizer/BioNTech

Pfizer-BioNTech COVID 19 VACCINE 2021 - 2022 FORMULA

INFORMATION SHEET (Pediatric Formulation)

Description

- The Pfizer/BioNTech vaccine provides protection against COVID 19. It will not prevent illness caused by other viruses. The vaccination will reduce the chance of you suffering from COVID-19 disease. Like all medicines, no vaccine is completely effective, and it takes a few weeks for your body to build up protection from the vaccine. Some people may still get COVID-19 despite having a vaccination, but this should lessen the severity of any infection.
- The vaccine cannot give you COVID-19 infection, and two doses will reduce your chance of becoming seriously ill. You will still need to follow the guidance in your workplace, including wearing the correct personal protection equipment and taking part in any screening programmes. Like all medicines, vaccines can cause side effects. Most of these are mild and short-term, and not everyone gets them.

COVID-19 vaccination requirements

- You will need two doses of the Pfizer/BioNTech COVID-19 vaccine for it to be fully effective; the two injections should be twelve weeks apart.
- You may not be protected until at least 7 days after the second dose of the vaccine.

Who should get vaccinated?

- All people five years of age and older should get the COVID 19 vaccine. This is especially important for people at higher risk for severe illness due to COVID 19 and their close contacts, including healthcare personnel and close contacts of children younger than twelve years old.
- Persons at higher risk of severe illness due to COVID 19 are: people (60 years of age and older), and people with certain health conditions such as heart, lung or kidney disease or a weakened immune system as COVID 19 may worsen existing medical conditions.
- Even if you have been sick with COVID 19 and have recovered, you are encouraged to receive the COVID 19 vaccine. Due to the life-threatening complications associated with COVID-19 and known cases of re-infection, you are encouraged to get the COVID-19 vaccine leaving at least four weeks from the time of infection.

Who should not be vaccinated or who should wait:

Some persons should not be vaccinated without first consulting a physician. They include:

- Children less than five years of age.
- Any person with a history of immediate onset of severe allergic reaction to a vaccine, medicine or food should not receive the Pfizer/BioNTech vaccine. A second dose of the Pfizer/BioNTech vaccine should not be given to those who have experienced immediate- onset of severe allergic reaction to the first dose of Pfizer/BioNTech vaccination.
- People who have a moderate or severe illness with a fever (more than 38°C) should wait to be vaccinated.

Possible side effects Common Problems:

- Soreness, redness or swelling of the vaccine site
- Headache
- Fatigue
- Chills
- Fever
- Muscle aches and pain
- Nausea.

If these occur, they usually begin soon after the vaccine and last one to two days.

Uncommon or rare problems:

- Discomfort or feeling ill
- Lymphadenopathy (swollen or enlarged lymph nodes)

Life-threatening allergic reactions from vaccines are very rare. When they do occur, it is usually within a few minutes to a few hours of receiving the vaccine.

Treatment of side effects

If you experience any side effects, please call the flu hotline at 1-800-534-8600 OR Text/WhatsApp 947-3077. If you have any difficulty breathing, call 911.

For more information

Talk to your healthcare provider; call your district health centre or refer to the question-and-answer brochure. You may also go online to the PHE website https://www.gov.uk

