

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/327918904>

A Connectomic Atlas of the Human Cerebrum—Chapter 10: Tractographic Description of the Superior Longitudinal Fasciculus

Article in *Operative Neurosurgery* · September 2018

DOI: 10.1093/ons/opy264

CITATIONS

5

READS

181

9 authors, including:



Andrew Conner

University of California, San Francisco

80 PUBLICATIONS 275 CITATIONS

[SEE PROFILE](#)



Robert G. Briggs

University of Southern California

57 PUBLICATIONS 210 CITATIONS

[SEE PROFILE](#)



Cordell Baker

University of Utah

40 PUBLICATIONS 162 CITATIONS

[SEE PROFILE](#)



Joshua Burks

University of Miami Miller School of Medicine

56 PUBLICATIONS 227 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Human brain networks and neurosurgery [View project](#)



Connectomic Atlas of the Human Cerebrum [View project](#)

A Connectomic Atlas of the Human Cerebrum—Chapter 10: Tractographic Description of the Superior Longitudinal Fasciculus

Andrew K. Conner, MD*

Robert G. Briggs, BS*

Meherzad Rahimi, BS*

Goksel Sali, MD*

Cordell M. Baker, MD*

Joshua D. Burks, MD*

Chad A. Glenn, MD*

James D. Battiste, MD, PhD[‡]

Michael E. Sughrue, MD*[§]

*Department of Neurosurgery, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma; [‡]Department of Neurology, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma; [§]Department of Neurosurgery, Prince of Wales Private Hospital, Sydney, Australia

Correspondence:

Michael E. Sughrue, MD,
Department of Neurosurgery,
Prince of Wales Private Hospital,
Level 7, Suite 3 Barker St.,
Randwick, NSW 2031, Australia
E-mail: sughruevs@gmail.com

Received, May 17, 2018.

Accepted, September 18, 2018.

Published Online, September 27, 2018.

Copyright © 2018 by the
Congress of Neurological Surgeons

The superior longitudinal fasciculus/arcuate white matter complex (SLF/AC) is the largest and most complex white matter tract of the human cerebrum with multiple inter-linked connections encompassing multiple cognitive functions such as language, attention, memory, emotion, and visuospatial function. However, little is known regarding the overall connectivity of this complex. Recently, the Human Connectome Project parcellated the human cortex into 180 distinct regions. Utilizing diffusion spectrum magnetic resonance imaging tractography coupled with the human cortex parcellation data presented earlier in this supplement, we aim to describe the macro-connectome of the SLF/AC in relation to the linked parcellations present within the human cortex. The purpose of this study is to present this information in an indexed, illustrated, and tractographically aided series of figures and tables for anatomic and clinical reference.

KEY WORDS: Anatomy, Cerebrum, Connectivity, DTI, Functional connectivity, Human, Parcellations

Operative Neurosurgery 00:S407–S422, 2018

DOI: 10.1093/ons/opy264

Having finished characterizing the structural and functional connectivity of all 180 cortical regions described in the Glasser study,¹ we now turn our attention to the specific tractographic fiber connections described in the first nine chapters of this supplement. Using these data, we define the sets of parcellations that integrate within the human brain to form eight large white matter tracts, including the superior longitudinal fasciculus (SLF), inferior longitudinal fasciculus, middle longitudinal fasciculus, inferior fronto-occipital fasciculus, frontal aslant tract, uncinate, cingulum, and vertical occipital fasciculus. We begin with the SLF that for our purposes includes the arcuate fasciculus (AF), making this the largest and most complex long-range fiber bundle in the brain.

The superior longitudinal fasciculus/arcuate white matter complex (SLF/AC) is the largest

white matter fiber grouping in the cerebrum, is present in each hemisphere, and is well known for its connections in the perisylvian, parietal, and occipital cortices.^{2,3} It is well described that the primary functional role associated with the SLF/AC is transmission of speech and language.⁴ It is also believed that the SLF plays a role in language transitioning (ie bilingualism).⁵ Other functional associations of the SLF/AC have also been described including crucial roles in attention, memory, emotion, language, visuospatial processing, and numerical cognition.^{6–8}

Although diffusion tensor imaging (DTI) studies have elucidated the structural anatomy of the SLF/AC, little is known about its various cortical terminations.⁹ Recently, the Human Connectome Project published parcellation data redefining the human cortex.¹ This provides a unique opportunity to elucidate the macro-connectome of the human cerebrum, in that high-resolution DTI tractography has been shown to accurately illustrate the anatomy and microstructure of the SLF/AC.¹⁰ In this study, we utilized diffusion spectrum imaging (DSI) tractography in conjunction with the Glasser parcellation scheme to illustrate the macro-connectivity of the parcellations

ABBREVIATIONS: AF, arcuate fasciculus; DSI, diffusion spectrum imaging; DTI, diffusion tensor imaging; IPL, inferior parietal lobule; MR, magnetic resonance; ROI, region of interest; SLF, superior longitudinal fasciculus; SLF/AC, superior longitudinal fasciculus/arcuate white matter complex

integrating within the confines of the SLF/AC. The purpose of this study is to present the structural connectivity of the SLF/AC in an indexed, illustrated, and tractographically aided series of figures and tables for anatomic and clinical reference.

METHODS


Identification of Relevant Cortical Regions



The parcellation data entries within the first nine chapters of this supplement were reviewed to determine the specific cortical regions with structural connectivity in the distribution of the SLF. These data were

tabulated, and connections between individual parcellations within the SLF were recorded. These results served as the basis for constructing simplified tractography maps of the SLF and performing deterministic tractography.


Deterministic Tractography

Publicly available imaging data from the Human Connectome Project was obtained for this study from the HCP database (<http://humanconnectome.org>, release Q3). Diffusion imaging with corresponding T1-weighted images from 10 healthy, unrelated controls were analyzed (subjects IDs: 100307, 103414, 105115, 110411,






111312, 113619, 115320, 117112, 118730, 118932). A multishell diffusion scheme was used, and the b-values were 990, 1985, and 1980 s/mm^2 . Each b-value was sampled in 90 directions. The in-plane resolution was 1.25 mm. The diffusion data were reconstructed using generalized q-sampling imaging with a diffusion sampling length ratio of 1.25.¹¹



We performed brain registration to montreal neurologic institute space, wherein imaging is warped to fit a standardized brain model comparison between subjects. Tractography was performed in DSI studio using a region of interest approach to initiate fiber tracking from a user-defined seed region. A two region of interest (ROI) approach was used to isolate tracts. Voxels within each ROI were automatically traced with a maximum angular threshold of 45°. When a voxel was approached with no tract direction or a direction change of greater than 45 degrees, the tract was halted. Tractography was stopped after reaching a maximum length of 800 mm. In some instances, exclusion ROIs were placed to exclude obvious spurious tracts that were not involved in the white matter pathway of interest. Tractographic results are shown only for regions of interest within the left cerebral hemisphere.

CONNECTIVITY OVERVIEW

Presented in Figure 1, we demonstrate the functionally relevant and anatomically connected cerebral parcellation data that integrates within the confines of the SLF/AC. In order to simplify this complex data set further, we divided the SLF/AC into three subunits based on the literature: SLF I, SLF II, and SLF III combined with the arcuate (Figures 2-4).¹² Pertinent examples of tractographically connected parcellations are represented for each subdivision (Figures 5-14). It should be noted that the figures and tables presented in this study do not imply directionality. Instead, supposed information transit is utilized as a simplified means



for connectivity description and reference. Given the variability between visual and descriptive renditions of the divisions of the SLF/AC, certain instances of overlap between parcellations are inevitable. The following subdivisions do not stand to concretely separate this tract. They are merely represented in this regard for ease of reference.

SLF Connectivity

Table lists the tractographically connected parcellations within the SLF/AF complex as a whole. Figure 1 illustrates the macro-connectome of these connections. Arrows are included for reference only. Directionality of information transit is not meant to be assumed or implied.





FIGURE 6. SLF connections from **A** and **B**, STSva and **C** and **D**, STSda to areas PSL and STV of the IPL. Connections are shown in the left cerebral hemisphere on T1-weighted MR images in the sagittal plane. All parcellations are identified with white arrows and corresponding labels. The SLF can be seen coursing posteriorly and superiorly within the subcortical white matter of the temporal before terminating in the perisylvian gray matter of the IPL.

SLF Connectivity

The SLF is classically divided into four distinct parts in human brains.¹² SLF I connects parts of the superior parietal lobule with secondary motor areas, including the supplementary motor area, in the superior frontal gyrus.¹³ Connectivity for

this subdivision is represented graphically in Figure 2. In short, connections in this region involve the medial complex of the SLF towards the medial interparietal sulcus, connecting areas underlying the superior frontal gyrus and the superior parietal lobule.



SLF II Connectivity

SLF II classically connects parts of the inferior parietal lobule (IPL) to the dorsolateral prefrontal cortex, including parts of the middle frontal gyrus.¹² Connectivity for this subdivision is represented graphically in Figure 3. In short, connections in this region involve the posterolateral complex of the SLF, inferior to

SLF I, connecting the parcellation of the middle frontal gyrus to the IPL.

SLF III/Arcuate Connectivity

SLF III classically runs within the frontal and parietal opercula, connecting parts of the IPL to the ventral prefrontal




FIGURE 8. A-C, SLF connections from area 44 in the inferior frontal gyrus to parts of temporal lobe. Connections are shown in the left cerebral hemisphere on T1-weighted MR images in the **A** and **B**, sagittal and **C**, axial planes. Area 44 exhibits structural connections via the SLF to temporal regions TGd, TE1a, TE1m, STSda, STSdp, and PHT in this subject brain. **D-F**, Area 44 also demonstrates structural connections to areas A5, A1, LBelt, and RI in the superior temporal gyrus and posterior insular cortex. Connections are shown in the left cerebral hemisphere on T1-weighted MR images in the **D** and **E**, sagittal, and **F**, axial planes. All parcellations are identified with white arrows and corresponding labels. The SLF can be seen coursing posteriorly before bending 90° to enter the subcortical white matter of the temporal lobe and terminate in the aforementioned parcellations.






FIGURE 9. SLF connections from parietal parcellation **A** and **B**, AIP to regions 6r and 43 of the inferior frontal gyrus. Connections are shown in the left cerebral hemisphere on T1-weighted MR images in the **A**, sagittal and **B**, axial planes. **C** and **D**, SLF connections from parietal parcellation LIPd to regions 55b and PEF of the frontal lobe. Connections are shown in the left cerebral hemisphere on T1-weighted MR images in the **C**, sagittal and **D**, axial planes. All parcellations are identified with white arrows and corresponding labels. The SLF can course inferiorly from the gray matter of the parietal lobe and gradually curves 90 degrees to terminate in different parts of the frontal lobe.



cortex, including parts of the inferior frontal gyrus.¹² Connectivity for this subdivision is represented graphically in Figure 4. In short, connections in this region involve the fronto-parietal operculum and its various connections to the insula. For simplicity, the arcuate complex is included in our depic-


tions of SLF III. The AF is classically considered the longest subdivision of the SLF, connecting the frontal operculum and preopercular parcellations to those in the temporal lobe.¹² Numerous connections to the perisylvian region are present in this scheme.



DISCUSSION

In this study, we describe the most detailed map to date regarding macro-connectivity of the SLF/AC and its relevant cerebral parcellations. It can be surmised from this

data that actionable future studies and surgical planning may be better outlined. Our data confirm the results of others,^{12,14-17} our descriptive study supports supposed connectivity within the confines of the human cerebrum. Anatomic models of the SLF/AC generally begin at the caudal portion



of the superior temporal gyrus, arch posteriorly and then caudally over the sylvian fissure to end within the frontal lobe.^{18,19}

Historically, the SLF/AC was thought to encompass a single entity and route of transmission void of multiple subunits.¹² However, recent evidence suggests the SLF can be divided into multiple subcomponents which can be described based on anatomic structures they connect.^{12,14-17} The SLF has also been divided into a superficial anterior segment connecting the supramarginal gyrus and superior temporal gyrus with the precentral gyrus and lateral frontal cortex, as well as a superficial posterior segment connecting the posterior portion of the middle temporal gyrus and supramarginal gyrus. Included within this paradigm is a long segment, otherwise called the AF, connecting the middle and inferior temporal gyri with the precentral, inferior, and middle frontal gyri.^{9,19}


From a purely structural standpoint, it is reasonable to hypothesize that the SLF/AC would play a critical role in language function given its known connections to the posterior language areas and the canonical Broca's area.²⁰ Studies regarding ideomotor apraxia discovered in the 1980s led to the idea that the SLF/AC is at least in part involved in speech articulation.²¹ Specifically, as illustrated during awake cortical and subcortical mapping studies, excellent correlation exists with the cortical boundary of the SLF and demonstrable speech function.²² Electrical stimulation of the SLF has also been shown

to elicit syntactic and phonemic disorders (ie, paraphasias) in the dominant cerebral hemisphere.²³ In addition, the language network has long been thought of as comprising a dorsal and ventral stream in which the dorsal pathway involves the SLF/AF complex.²⁴ Damage to the SLF/AF is associated with nonfluent variants of primary progressive aphasia, again suggesting that the SLF is critical to normal language function.²⁴ Finally, the dorsal phonological system, with deficits characterized by phonemic paraphasia, has been demonstrated to be integrated within the SLF, specifically localizing to the area of the IPL with connections between the supramarginal gyrus and ventral premotor cortex.²⁵

On the nondominant side, the SLF is thought to play a role in visuospatial awareness as well as attentional selection of sensory content. This is further reinforced secondary to studies demonstrating hemi-neglect subsequent to damage to the inferior parietal lobe and SLF as it runs in this region.²⁶⁻³² The SLF may also play an important role in the modulation of audio-spatial information.⁷

CONCLUSION

The SLF/AC is an incredibly complex white matter tract connecting multiple regions of the human cerebrum. It is critical



in the transmission and processing of multiple types of data for the proper execution of cognitive tasks such as language, attention, memory, emotion, and visuospatial function. Further, sub-tract-guided functional and anatomic studies are needed

to enhance our understanding of the functional connectivity of the human cerebrum. Our tractographic map of this white matter bundle can serve as a reference point for these future studies.




FIGURE 14. SLF connections from inferior frontal sulcus area PFm. Area PFm demonstrates structural connections to **A** and **B**, areas 2, 4, 55b, and 8Av anteriorly and **C** and **D**, areas STSvp, TE1m, TE1p, and PHT inferiorly. These connections are shown in the left cerebral hemisphere on T1-weighted MR images in the **A** and **C**, sagittal and **B** and **D**, axial planes, respectively. All parcellations are identified with white arrows and corresponding labels.

TABLE. Regions Integrating Within the SLF

Original parcellation	Terminations
2	IP1
	IP2
	PFm
3a	6a
3b	IP1
	IP2
6r	PSL
	RI
	STV
	TE1a
	TE2a
	TE2p
	TPOJ1
6v	FST
	IP1
	IP2
	PH
	PHT
	TE2a
	TE2p
	TPOJ2
8AV	2
	6a
	7PC
	MIP
8C	PH
	PHT
	TE2a
44	A1
	A5
	LBelt
	PHT
	RI
	STSdp
	STSva
	TE1a
	TE1m
	TGd
45	A4
	PBelt
	TE1m
	TE1p
46	TF
55b	IP1
	IP2
	PHT
	TGd
AIP	6r
	43
FEF	IP1
	IP2
FOP1	TE1a
FOP4	A4
	A5
	STSdp
	TE1a
	TGd

TABLE. Continued.

Original parcellation	Terminations
FOP5	A4
	LBelt
	PBelt
	RI
IFJa	PHT
	STSdp
	STSva
	STSvp
	TE1a
	TE1m
	TE2a
	TPOJ1
	TPOJ2
IFJp	FST
	PHT
	STSdp
	STSvp
	TE1m
	TE2a
	TPOJ1
	TPOJ2
IFSa	4
	TE2a
	TF
IFSdp	STSdp
	TE1a
	TE1m
	TE2a
LIPd	55b
	PEF
MI	A4
	A5
	RI
p9-46v	TE2a
PEF	IP1
	IP2
PF	1
	3a
	4
	6v
	43
	OP4
	STSva
	TE1a
	TE2a
PFcm	43
	TE1a
	TE2a
PFm	1
	2
	3a
	4
	8AV
	8BM
	8C
	43
	55b

TABLE. Continued.

Original parcellation	Terminations
	STSva
	STSvp
	TE1a
	TE1m
	TE1p
PFT	4
	6r
	43
	OP4
PGs	FEF
	FST
	IFJa
	IFJp
	IFSp
	PHT
	TPOJ2
STSda	FOP1
	FOP3
	FOP4
	STV
STSdp	IFSp
	STV
	PSL
STSva	6r
	43
	PFcm
	PSL
	STV
STSvp	6r
	FOP2
	FOP3
	PSL
	STV
TE1a	FOP3
	PSL
	STV

Disclosures

Synaptive Medical assisted in the funding of all 18 chapters of this supplement. No other funding sources were utilized in the production or submission of this work.

REFERENCES

1. Glasser MF, Coalson TS, Robinson EC, et al. A multi-modal parcellation of human cerebral cortex. *Nature*. 2016;536(7615):171-178.
2. Jellison BJ, Field AS, Medow J, Lazar M, Salamat MS, Alexander AL. Diffusion tensor imaging of cerebral white matter: a pictorial review of physics, fiber tract anatomy, and tumor imaging patterns. *AJNR Am J Neuroradiol*. 2004;25(3):356-369.
3. Schmahmann JD, Smith EE, Eichler FS, Filley CM. Cerebral white matter: neuroanatomy, clinical neurology, and neurobehavioral correlates. *Ann N Y Acad Sci*. 2008;1142:266-309.
4. Axer H, Klingner CM, Prescher A. Fiber anatomy of dorsal and ventral language streams. *Brain Lang*. 2013;127(2):192-204.
5. Moritz-Gasser S, Duffau H. Cognitive processes and neural basis of language switching: proposal of a new model. *Neuroreport*. 2009;20(18):1577-1580.
6. Mesulam MM. From sensation to cognition. *Brain*. 1998;121 (6):1013-1052.
7. Petrides M, Pandya DN. Comparative cytoarchitectonic analysis of the human and the macaque ventrolateral prefrontal cortex and corticocortical connection patterns in the monkey. *Eur J Neurosci*. 2002;16(2):291-310.
8. Moeller K, Willmes K, Klein E. A review on functional and structural brain connectivity in numerical cognition. *Front Hum Neurosci*. 2015;9:227.
9. Martino J, De Witt Hamer PC, Berger MS, et al. Analysis of the subcomponents and cortical terminations of the perisylvian superior longitudinal fasciculus: a fiber dissection and DTI tractography study. *Brain Struct Funct*. 2013;218(1):105-121.
10. Kamali A, Flanders AE, Brody J, Hunter JV, Hasan KM. Tracing superior longitudinal fasciculus connectivity in the human brain using high resolution diffusion tensor tractography. *Brain Struct Funct*. 2014;219(1):269-281.
11. Yeh FC, Wedeen VJ, Tseng WY. Generalized q-sampling imaging. *IEEE Trans Med Imaging*. 2010;29(9):1626-1635.
12. Makris N, Kennedy DN, McNerney S, et al. Segmentation of subcomponents within the superior longitudinal fascicle in humans: a quantitative, in vivo, DT-MRI study. *Cereb Cortex*. 2005;15(6):854-869.
13. Jang SH, Hong JH. The anatomical characteristics of superior longitudinal fasciculus I in human brain: diffusion tensor tractography study. *Neurosci Lett*. 2012;506(1):146-148.
14. Catani M, Jones DK. Perisylvian language networks of the human brain. *Ann Neurol*. 2005;57(1):8-16.
15. Frey S, Campbell JS, Pike GB, Petrides M. Dissociating the human language pathways with high angular resolution diffusion fiber tractography. *J Neurosci*. 2008;28(45):11435-11444.
16. Makris N, Papadimitriou GM, Kaiser JR, Sorg S, Kennedy DN, Pandya DN. Delineation of the middle longitudinal fascicle in humans: a quantitative, in vivo, DT-MRI study. *Cereb Cortex*. 2009;19(4):777-785.
17. Zhang Y, Zhang J, Oishi K, et al. Atlas-guided tract reconstruction for automated and comprehensive examination of the white matter anatomy. *Neuroimage*. 2010;52(4):1289-1301.
18. Christidi F, Karavasilis E, Samiotis K, Bisdas S, Papanikolaou N. Fiber tracking: A qualitative and quantitative comparison between four different software tools on the reconstruction of major white matter tracts. *Eur J Radiol Open*. 2016;3: 153-161.
19. Potapov AA, Goryainov SA, Zhukov VY, et al. [The long-associative pathway of the white matter: modern view from the perspective of neuroscience]. *Zh Vopr Neurokhir Im N N Burdenko*. 2014;78(5):66-77; discussion 77.
20. Bernal B, Altman N. The connectivity of the superior longitudinal fasciculus: a tractography DTI study. *Magn Reson Imaging*. 2010;28(2):217-225.
21. De Renzi E. [Agnosia]. *Recenti Prog Med*. 1989;80(12):633-637.
22. Davtian M, Ulmer JL, Mueller WM, Gaggl W, Mulane MP, Krouwer HG. The superior longitudinal fasciculus and speech arrest. *J Comput Assist Tomogr*. 2008;32(3):410-414.
23. Wilson SM, Galantucci S, Tartaglia MC, Gorno-Tempini ML. The neural basis of syntactic deficits in primary progressive aphasia. *Brain Lang*. 2012;122(3): 190-198.
24. Agosta F, Galantucci S, Canu E, et al. Disruption of structural connectivity along the dorsal and ventral language pathways in patients with nonfluent and semantic variant primary progressive aphasia: a DT MRI study and a literature review. *Brain Lang*. 2013;127(2):157-166.
25. Maldonado IL, Moritz-Gasser S, Duffau H. Does the left superior longitudinal fascicle subserve language semantics? A brain electrostimulation study. *Brain Struct Funct*. 2011;216(3):263-274.
26. Hoeft F, Barnea-Goraly N, Haas BW, et al. More is not always better: increased fractional anisotropy of superior longitudinal fasciculus associated with poor visuospatial abilities in Williams syndrome. *J Neurosci*. 2007;27(44): 11960-11965.
27. Naito E, Amemiya K, Morita T. [Parietal cortices and body information]. *Brain Nerve*. 2016;68(11):1313-1320.
28. Ptak R. The frontoparietal attention network of the human brain. *Neuroscientist*. 2012;18(5):502-515.
29. Chechlacz M, Rotshstein P, Hansen PC, Deb S, Riddoch MJ, Humphreys GW. The central role of the temporo-parietal junction and the superior longitudinal

- fasciculus in supporting multi-item competition: evidence from lesion-symptom mapping of extinction. *Cortex*. 2013;49(2):487-506.
30. Mesulam MM. A cortical network for directed attention and unilateral neglect. *Ann Neurol*. 1981;10(4):309-325.
 31. Shinoura N, Suzuki Y, Yamada R, Tabei Y, Saito K, Yagi K. Damage to the right superior longitudinal fasciculus in the inferior parietal lobe plays a role in spatial neglect. *Neuropsychologia*. 2009;47(12):2600-2603.
 32. Posner MI, Walker JA, Friedrich FJ, Rafal RD. Effects of parietal injury on covert orienting of attention. *J Neurosci*. 1984;4(7):1863-1874.

Acknowledgments

Data were provided [in part] by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University. We would also like to thank Brad Fernald, Haley Harris, and Alicia McNeely of Synaptive Medical for their assistance in constructing the network figures for Chapter 18 and for coordinating the completion and submission of this supplement.