

## **DOSING FOR INFANTS & KIDS |** FOR US HEALTHCARE PROFESSIONALS ONLY

**DOSE:** Find the right dose on the chart below. If possible, use weight to dose; otherwise, use age. Repeat every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS. Do NOT use with any other product containing acetaminophen.

	Weight	Age	Dose	
Infants' Pain & Fever	6 - 11 lbs.	0 - 3 months	1.25 mL	# <del>u</del> b
Acetaminophen 160 mg per 5mL Oral Suspension *Syringe included	12 - 17 lbs.	4 - 11 months	2.5 mL	
Genexa	18 - 23 lbs.	12 - 23 months	3.75 mL	#IIID
Infants' Pain & Fever	24 - 35 lbs.	2 - 3 years	5 mL	Ħ
Make and this Specie Market by Macro this common energies, a solition of sens, or yourseless apple in 2 mars - loan Intrinsic feature, by (2008) 2 FL OZ (56 mil.)				

American String as to, American String as to, Fire Nation - Food Paleon  American String as to the Control of the Control  American String as to the Control  American String as the Control  American String a				
Kids' Pain & Fever	24 - 35 lbs.	2 - 3 years	5 mL	
Acetaminophen 160 mg per 5mL Oral Suspension Ages 2-11	36 - 47 lbs.	4 - 5 years	7.5 mL	
*Dosing cup included Use product only as directed  Genexa  **Autobate **Autob	48 - 59 lbs.	6 - 8 years	10 mL	
	60 - 71 lbs.	9 - 10 years	12.5 mL	
	72 - 95 lbs.	11 years	15 mL	

# Infants' Pain & Fever Kids' Pain & Fever

Acetaminophen 160 mg per 5mL Oral Suspension



Ibuprofen-free

**Aspirin-free** 

**Alcohol-free** 

**No Common Allergens** 

**No Lactose** 

**No Parabens** 

**No Artificial Dyes** 

**No Artificial Flavors** 

**No Artificial Preservatives** 



Made with Organic Blueberry Flavor

## REMIND PARENTS AND CAREGIVERS ON IMPORTANT INSTRUCTIONS FOR PROPER USE:

- 1. Please refer to outer package for complete labeling and warning information.
- 2. **DO NOT USE** with any other product containing acetaminophen.
- 3. DO NOT give more than directed.
- 4. SHAKE WELL BEFORE USING.
- 5. Use only the measuring device provided with the product to measure the proper amount of medicine; do not use any other device.
- 6. DO NOT GIVE MORE THAN 5 TIMES IN 24 HOURS.
- 7. Replace cap tightly to maintain child resistance.
- 8. Store between 20-25°C (68-77°F).











## Genexa dosing for kids | for us healthcare professionals only

		Age	Dose	
Genexa  REAL MEDICINE  BASE OF THE STATE OF	Kids' Allergy Diphenhydramine HCl Liquid Ages 6+ *Dosing cup included Use product only as directed	adults and children 12 years of age and over	10 – 20 mL	
		children 6 to under 12 years	5 – 10 mL	
		children 2 to under 6 years	do not use unless directed by a doctor	
		children under 2 years	do not use	
				_

Kids' Cough & Chest Conges  Dextromethor Guaifenesin Oral Suspension  Ages 4-11 *Dosing cup include  *Dosing cup i	Kids' Cough & Chest	children 6 to under 12 years	5 mL – 10 mL every 4 hours
	Congestion  Dextromethorphan HBr / Guaifenesin Oral Suspension	children 4 to under 6 years	2.5 mL – 5 mL every 4 hours
	Ages 4-11 *Dosing cup included Use product only as directed	children under 4 years	do not use

Age



Kids' Tummy Relief Calcium Carbonate Chewable Tablets Ages 2-11

Weight	Age	Dose	
over 48 lbs.	6-11 years	chew 2 tablets as symptoms occur, not to exceed 6 tablets in 24 hours	(1) (1)
24-47 lbs.	2-5 years	chew 1 tablet as symptoms occur, not to exceed 3 tablets in 24 hours	<b>(</b> )
under 24 lbs.	under 2 years	ask a doctor	



Dose



Kids' Immune Support **Dietary Supplement** 

Ages 2+

## For Daily Maintenance:

- · children 2-3 years of age: take 1 tsp (5mL) daily
- · children 4 years of age and older: take 2 tsp (10mL) daily

### **For Intensive Support:**

- · children 2-3 years of age: take 1 tsp (5mL) four times daily
- · children 4 years of age and older: take 2 tsp (10mL) four times daily



Kids' Honey Cough Syrup **Dietary Supplement** 

- · children 1-3 years of age: take one ½ tsp (2.5 mL) four times daily as needed for 5-7 days
- · children 4 years of age and older: take 1 tsp (5 mL) four times daily as needed for 5-7 days